



Research article

## Preliminary phytochemical screening of raw and thermally processed palmyra palm (*Borassus flabellifer* linn.) fruit pulp

P.Saranya, T. Poongodi Vijayakumar\*

Department of Food Science and Nutrition, Periyar University, Periyar Palkalai Nagar, Salem, Tamilnadu, India.

### Abstract

Palmyra palm (*Borassus flabellifer* Linn.) can be found in tropical countries such as Thailand, Malaysia, Indonesia, India, Myanmar, Sri Lanka and Cambodia. The palmyra palm (*Borassus flabellifer* Linn.) fruit is one of the less known tropical fruits. The major reasons for the underutilization of these fruits are separation of the pulp from the fiber and its bitter taste. The fruits are stomachic, sedative, laxative and aphrodisiac in nature. The current study was conducted to screen the different phytochemicals including alkaloids, flavonoids, terpenes, glycosides, saponins, phenolics, tannin, carbohydrate, protein and steroids and sterols from aqueous and methanolic extracts of raw palmyra palm fruit pulp (RPF) and thermally processed palmyra palm fruit pulp (PPF). The result revealed that both the RPF and PPF contain all the phytochemicals except protein. The terpenes, glycosides, carbohydrate and steroids and sterols are present in appreciable amount and proteins are absent in all the extraction of both samples. Alkaloids were present appreciably in the aqueous extraction of RPF and present fewer amounts in PPF. But methanol extraction of both RPF and PPF contain alkaloids strongly. The flavonoids, phenols and tannins were present in both the extraction of RPF and PPF. Glycosides were found in the aqueous extraction but present appreciable amount in methanol extraction of RPF and PPF. On processing alkaloids were reduced in the aqueous extraction of PPF and saponin in both the extraction of RPF and PPF. Therefore no major changes in the presence of phytochemical components due to heat processing were found in this current study.

**Key words:** Raw palm fruit pulp (RPF), processed palm fruit pulp (PPF), aqueous extraction, methanol extraction, phytochemical screening.

**\*Corresponding Author: T. Poongodi Vijayakumar**, Professor and Head, Department of Food Science and Nutrition, Periyar University, Periyar Palkalai Nagar, Salem- 636 011, TamilNadu, India.

### 1. Introduction

Palmyra palm (*Borassus flabellifer* Linn.) can be found in tropical countries such as

Thailand, Malaysia, Indonesia, India, Myanmar, Sri Lanka and Cambodia [1].

The palmyra palm is known simply as palmyra, which is based on the Portuguese palmeira, the tree was named originally for the resemblance of the leaf to the palm of a hand. Other names are toddy palm, wine palm, Cambodian palm and botanically known as *Borassus flabellifer* L. *Borassus* is from a Greek word describing the leathery covering of the fruit, Flabellifer means "fan-bearer" [2]. The palm belongs to the family *Arecaceae*, subfamily *Borassoideae* and genus *Borassus*. The palmyrah palms are slow-growing dioecious perennials and have no distinguishing features to identify the sex until flowering. The palm commences flowering only after 12–15 years of maturity. Inflorescences are interfoliar; the male inflorescence has stout terete branches, while the female inflorescence is more sparingly branched. Fruits are semi-globose to globose and deep brown to black when ripe [3]. The palmyra palm (*Borassus flabellifer* Linn.) fruit is one of the less known tropical fruits. The coconut-like fruits are three-sided when young, becoming rounded or more or less oval, 12-15 cm wide and capped at the base with overlapping sepals. The outer covering is smooth, thin, leathery and brown, turning nearly black after harvest. Inside is a juicy mass of long, tough, coarse, white fibers coated with yellow or orange pulp. The pulp of mature fruits is sucked directly from the wiry fibers of roasted, peeled fruits. It is also extracted to prepare a product called punatoo in Ceylon. It is eaten alone or with the starch from the palmyra seedlings. The fresh pulp is reportedly rich in vitamins A and C [4]. Jayaratnam (1986) isolated tetraglycoside of the steroid and later the bitter compound called flabelliferrins, which are steroidal saponins and this bitter principle in *Borassus* was identified

by Jansz *et al.* (1994) [5,6]. The major reasons for the underutilization of these fruits are separation of the pulp from the fiber and its bitter taste. Debittering is an important process for the utilization of the PFP. Heating the palmyra palm fruit over hot coals traditionally or using heat stable enzymes scientifically is reduced the bitterness [6,7].

The different parts of the plant are used for the various ailments like secondary syphilis, antiperiodic, heart burns, liver and spleen enlargement [8]. The palmyrah plant has been used traditionally as a stimulant, anti-leprotic, diuretic, antiphlogistic. The fruits are stomachic, sedative, laxative and aphrodisiac in nature. The roots and juice of the plant are useful in inflammatory reactions [9,10,11].

Phytochemicals are the bioactive, non-nutrient, and naturally occurring plant compounds found in fruits, vegetables, and whole grains. They can be categorized into various groups, i.e., polyphenols, organo-sulfur compounds, carotenoids, alkaloids, and nitrogen-containing compounds [12]. The preliminary phytochemical screening tests may be useful in the detection of the bioactive principles and subsequently may lead to the drug discovery and development. Further, these tests facilitate their quantitative estimation and qualitative separation of pharmacologically active chemical compounds [13]. There is the potential use of these tropical fruit pulps and their by-products to isolate specific phytochemicals for application in nutraceutical supplements, dietary additives, new food and pharmaceutical products, contributing to the recovery of agro-industrial process waste, with major industrial, economic and environmental impact. Therefore, the identification and

quantification of phytochemicals in the pulps and by-products of tropical fruits are of utmost importance to substantiate their potential health benefits in human nutrition [14].

Hence the present study was aimed at qualitative phytochemical screening from the raw palmyra palm fruit pulp (RPF) and processed palmyra palm fruit pulp (PPF) for further quantification of bioactive components.

## **2. Materials and method**

### **Sample collection**

The fully ripened palm fruits (50 samples) with distinct flavors and dispersed easily from the crown were randomly picked from the tree with the help of palm tree climber.

### **Pulp Extraction**

The ripe palmyra fruits were thoroughly washed, the outer black skin was peeled off; fibrous part with fruit pulp were separated from the seed and extracted by home scale juicer. One part of the fresh pulp was filled in sterilized glass bottles and stored in the refrigerator and the rest were heated at 90°C for 15 minutes, potassium meta bisulphate (0.14 g/Kg of pulp) was added as a preservative packed into sterilized glass bottles and finally the filled bottles were pasteurized (80°C for 20 min) for further use.

### **Aqueous extract**

The aqueous extraction was done by the cold maceration method described by Anowi (2012) with slight modifications in palmyra palm fruit pulp samples (RPF and PPF) separately [15]. 20 g of fresh fruit pulp was added to 200ml of water and allowed to stand for twenty four (24) hours. It was made into slurry by blending

with another 100ml of water and then filtered through Whatman no. 1 filter paper and the extract was directly used for qualitative tests.

### **Methanol extraction**

The alcoholic extraction method according to Cowan (1999) was used for methanol extraction of palmyra palm fruit pulp samples (RPF and PPF) separately [16]. About 20 g of the prepared palmyra palm fruit pulp samples were soaked in 200ml of methanol for extended periods (about 24 hours). The slurry was then filtered through Whatman no. 1 filter paper and washed with 50ml of methanol, after which it was used directly for the qualitative phytochemical screening.

### **Qualitative analysis of Phytochemical screening**

The following the methods of Horbone, (1973) and Trease and Evans (1989) were used in the phytochemical screening of aqueous and methanol extraction of palmyra palm fruit pulp (RPF and PPF) [17,18].

### **Test for Alkaloids**

Crude extract was mixed with 2 ml of 1% HCl and heated gently. Mayer's and Wagner's reagent were added to the mixture. Appearance of cream color precipitates with Mayer's reagent and appearance of reddish brown precipitates with Wagner's reagent indicates the presence of alkaloids.

### **Test for Flavonoids**

Crude extract was mixed with 5ml of dilute ammonia followed by the addition of concentrated sulphuric acid. A yellow coloration observed in the extract indicated the presence of flavonoids. The yellow coloration disappears on standing.

#### **Test for Terpenoids (Salkowski test)**

Crude extract was mixed with 2ml of chloroform and concentrated sulphuric acid was added sideways. A reddish brown coloration at the interface indicates the presence of terpenoids.

#### **Test for Cardiac glycosides (Keller-Killani test)**

Crude extract was mixed with 2ml of glacial acetic acid containing 1-2 drops of 2% solution of  $\text{FeCl}_3$ . The mixture was then poured into another test tube containing 2 ml of concentrated sulphuric acid. A brown ring at the interphase indicates the presence of cardiac glycosides.

#### **Test for Saponins**

Crude extract was mixed with 5ml of distilled water in a test tube and shaken vigorously. The formation of stable foam was taken as an indication for the presence of saponins.

#### **Test for Phenols**

Crude extract was mixed with a few drops of 10 % solution of lead acetate. White precipitate indicates the presence of phenols.

#### **Test for Tannins**

Crude extract was mixed with 2 ml of neutral  $\text{FeCl}_3$ . A dark green coloration indicates the presence of tannins.

#### **Test for Carbohydrates**

Crude extract was mixed with a few drops of  $\alpha$  naphthol solution in alcohol and concentrated sulphuric acid was added from the side of the test tube. Violet ring formed at the junction of two liquids showed the presence of carbohydrate.

#### **Test for Proteins**

Crude extract was mixed with 2 ml of Biuret reagent. The violet color indicates the presence of proteins.

#### **Test of Steroids and sterols**

The extract was dissolved in 2 ml of chloroform and equal volume of concentrated sulphuric acid was added along the sides of the test tube. The upper layer turns red and lower layer turns yellow with green fluorescence, indicating the presence of the steroids and sterols compound, in the extract.

#### **Results and Discussion**

The aqueous and methanolic extracts of RPF and PPF were screened for different phytochemicals including alkaloids, flavonoids, terpenes, glycosides, saponins, phenolic, tannin, carbohydrate, protein and steroids and sterols. The results of both aqueous and methanol extraction for phytochemical tests (Table 1) indicated that the RPF and PPF contain alkaloids, flavonoids, saponins, phenolics, tannins appreciably. The strong presence of alkaloids, flavonoids, phenols and tannins were evidence by the formation of strong colour and precipitate. The presence of terpenes, glycosides, carbohydrate and steroids, sterols and absence of proteins were also noticed in all the extraction of both samples. Alkaloids were present appreciably in the aqueous extraction of RPF and present fewer amounts in PPF. But methanol extraction of both RPF and PPF contain alkaloids strongly. The flavonoids, phenols and tannins were present in both the extraction of RPF and PPF. Glycosides were found in the aqueous extraction, but present appreciable amount in methanol extraction of RPF and PPF. The ethanolic, chloroform and aqueous extracts of fresh palmyrah fruit

pulp and sun dried fruit pulp (Pannatu) were studied and the extracts contain steroids, triterpenoids, carbohydrates, saponin, flavonoids and proteins in varied amounts in the ethanolic and water extracts which is similar to this current study except for protein. The chloroform extract showed negative results for all tested compounds except for carbohydrate in the same [19]. Glycosides, alkaloids and tannins were not observed in any of the extracts but all these were present in the both aqueous and methanolic extract of PFP samples in the current study. Another study showed that the ethanolic extract of PFP powder indicated the presence of saponins, flavonoids, glycosides and phenolic compounds [20]. In contrary, the alkaloids and tannins were absent in the same. The differences might be due to the varietal changes of fruit and different solvents used for the extraction.

On processing alkaloids were reduced only in the aqueous extraction of PFP. The saponin was also decreased in both aqueous and methanolic extraction of RFP and PFP. The level of phytochemicals in vegetable and fruit processing decreases exponentially with a linear increase in blanching and boiling time [21]. Processing of fruit or vegetables can result in a significant reduction in phytochemical content. There are no changes in the presence of flavanoids, phenols and tannin in the raw PFP (RFP) after thermal processing (PFP). The results from phytochemical screening of bitter garden egg (*Solanum incanum*) fruit indicated that there is no much difference in raw and heat processed samples for 15 minutes [22] Glycosides are strongly present in the methanolic extraction than aqueous extraction which indicates the different solvents of different polarity is

used to separate compounds based on their solubility in the extraction solvent [23].

Alkaloids are one of the largest groups of phytochemicals that have led to the invention of powerful pain killer medications [24]. They are also involved in protective function in animals and are used as medicine, especially the steroidal alkaloids [25]. Steroids are known to be an important Cardio tonic activities posse's antimicrobial property and also used in herbal medicines and cosmetics. The presences of cardiac glycosides are known to play a major role in heart muscles by inhibiting Na<sup>+</sup> and K<sup>+</sup> pump that increases the availability of sodium ions and calcium ions to heart muscles which improves cardiac output and reduce heart distension. Thus are used in the treatment of congestive heart failure and cardiac arrhythmia [26]. Flavanoids and tannin are the group of phenolic compounds that act as primary antioxidants and posse's antimicrobial, anti-inflammatory, anti-allergic, anticancer, anti-neoplastic activity, and for the treatment of intestinal disorders [27]. In Ayurveda, formulations based on tannin-rich plants have been used for the treatment of diseases like leucorrhoea, rhinorrhoea and diarrhea [28] Saponins which act as bioactive antibacterial agents in plants are also used to treat hypercholesterolemia, hyperglycemia and obesity [29]. Saponins are also important therapeutically as they are shown to have hypolipidemic and anticancer activity. Saponins are also necessary for activity of cardiac glycosides [28]. Fruits that have tannins as their major components are astringent in nature. They are used in treating intestinal disorders such as diarrhea and dysentery as in [30, 31].

**Table 1. Phytochemical screening of aqueous and methanol extract of RPF and PPF**

Phytochemicals	Aqueous Extraction		Methanol Extraction	
	RPF	PPF	RPF	PPF
Alkaloids	++	+	++	++
Flavonoids	++	++	++	++
Terpenoids	+	+	+	+
Cardiac Glycosides	+	+	++	++
Saponins	++	+	++	+
Phenolics	++	++	++	++
Tannin	++	++	++	++
Carbohydrate	+	+	+	+
Protein	-	-	-	-
Steroids and Sterols	+	+	+	+

++ indicates strong presence, + indicates presence - indicates absence

### Conclusion

The preliminary phytochemical screening is the fundamental step for quantitative estimation of bioactive chemical constituents present in the fruit pulp. There was no major alteration after thermal processing in the qualitative phytochemical compounds of the palmyra palm fruit pulp. Thus the phytochemical compounds present in the raw and processed palmyra palm fruit pulp can be extracted for the further drug development.

### References

1. Naknean P, Meenune M and Roudaut G: Characterization of palm sap harvested in Songkhla province, Southern Thailand. *International Food Research Journal* 2010; 17: 977-986.
2. Small E: *Top 100 Exotic Food Plants*. CRC Press, Boca Raton 2011; 443-448.
3. Ramachandran VS, Swarupnandan K and Renuka C: A Traditional Irrigation System Using Palmyra Palm (*Borassus flabellifer*) in Kerala, India. *Palms* 2004; 48(4): 175-181.
4. Morton JF: Notes on distribution, propagation and products of *Borassus Palms* (Arecaceae). *Economic Botany* 1988; 42(3): 420-441.
5. Jansz ER, Nikawala JK, Goonaratne J and Theivendirarajah K: Studies on the bitter principle and debittering of palmyrah fruit pulp. *Journal of Science of Food and Agriculture* 1994; 65:185-189.
6. Jeyaratnam M: Studies on the chemistry and biochemistry of palmyrah products. M.Phil. Thesis, University of Jaffna, Sri Lanka, 1986; 1-200.
7. Ariyasena DD: The Diversity, Bioactivity and structural studies on the flabelliferins of palmyrah (*Borassus flabellifer* L.) fruit pulp. M.Phil. Thesis, University of Sri Jayewardenepura, Sri Lanka, 2002; 12-26.
8. Sandhya S, Sudhakar K, Banji D and Rao KNV: Pharmacognostical standardization of *Borassus Flabellifer* root. *Annals of Biological Research* 2010; 1 (4):85-94.
9. Nadkarni KM: *Indian Materia Medica*, 3<sup>rd</sup> Ed. Popular Book Depot, Bombay, India 1954; 4:2571-2575.

10. Vaidyaratnam PS: Varier's Arya Vaidya Sala - Indian Medicinal Plants A Compendium of 500 species. Orient Longman, Chennai, India 1994; 4:293-296.
11. Kapoor LD: Handbook of Ayurvedic medicinal plants. Herbal reference library, CRC Press, USA, Florida 2000; 82.
12. Lampe JW and Chang JL: Interindividual differences in phytochemical metabolism and disposition. Seminars in cancer biology 2007; 17(5): 347-353.
13. Varadarajan P, Rathinaswamy G and Asirvatham D: Antimicrobial properties and phytochemical constituents of Rheo discolor. Ethnobotanical Leaflet 2008; 12: 841-845.
14. Ayala-Zavala, Rosas-Dominguez C, Vega-Vega V and Gonzalez-Aguilar GA: Antioxidant enrichment and antimicrobial protection of fresh-cut fruits using their own byproducts: Looking for integral exploitation. Journal of Food Science 2010; 25 (8):75-181.
15. Anowi CF, Nnabuife CC, Mbah CJ and Onyekaba TC: Antimicrobial Properties of the Methanolic Extract of the Leaves of *Nauclea latifolia*. International Journal of Drug Research and Technology 2012; 2: 45-55.
16. Cowan MM: Plant Products and Antimicrobial Agents. American Society for Microbiology 1999; 12: 564-582.
17. Horbone JB: In: Phytochemical methods, 2<sup>nd</sup> Ed. Chapman and Hall, Newyork 1984.
18. Trease GE and Evans WC: Pharmacognosy, 11<sup>th</sup> Ed. Bailliere Tindall, London 1989; 45-50.
19. Manoharan NA, Mahilrajan S and Sri Vijeindran S: Preliminary phytochemical screening of extracts of palmyrah fruit pulp. Proceedings of the Peradeniya Univ. International Research Sessions, Sri Lanka, 2014; 18; 545.
20. Vijayakumari B, Vengaiah PC and Kiranmayi P: Qualitative phytochemical screening, GC-MS analysis and antibacterial activity of palmyra fruit pulp (*Borassus flabellifer* L.). International Journal of Pharma and Bio Sciences 2015; 6(2):430 - 435.
21. Tiwari U and Cummins E: Factors influencing levels of phytochemicals in selected fruit and vegetables during pre- and post-harvest food processing operations. Food Research International 2013; 50:497-506.
22. Auta R, James SA, Auta T and Sofa EM: Nutritive Value and Phytochemical Composition of *Solanum incanum* (*Bitter garden* Egg). Science World Journal 2011; 6(3):5-6.
23. Rashmi S and Rajkumar HG: Preliminary phytochemical screening of different solvent extracts of lichens from Kodagu district, Karnataka. Journal of Pharmacognosy and Phytochemistry 2014; 3(4):209-212.
24. Kam PCA and Liew: Traditional Chinese Herbal medicine and anesthesia. Anesthesia 2002; 57 (11):1083-1089.
25. Lata N and Dubey V: Preliminary phytochemical screening of *Eichhornia crassipes*: the world's worst aquatic Weed. Journal of Pharmacy Research 2010; 3:1240-1242.
26. Schneider G and Wolfling J: Synthetic Cardenolides and related compounds. Current Organic Chemistry 2004; 8:14.
27. Rievere C, Nguyen JHV, Pieters L, Dejaegher B, Heyden YV, Minh CV and Quetin-Leclercq J: Polyphenols isolated from antiradical extracts of *Mallotus metcalfeanus*. Phytochemistry 2009; 70:86-94.
28. Doughari JH: Phytochemicals - Extraction Methods, Basic Structures and Mode of Action as Potential Chemotherapeutic Agents, Publisher InTech, Croatia (European Union) 2012; 1-32.
29. Mohanta TK, Patra JK, Rath SK, Pal DK and Thatoi HN: Evaluation of antimicrobial activity and phytochemical screening of oils and nuts of *Semicarpus anacardium* L. f. Sci. Res. 2007 Essay., 2(11):486-490.
30. Chrinius H, Gary GY, Gideon AS, Asabe MM and Abel SA: Phytochemical and antimicrobial screening of methanol and

- aqueous extracts of *Agave sisalana*. *Acta Poloniae Pharmaceutica-Drug Research* 2011; 68 (4):535-539.
31. Nisha Y, Gunjan T, Deepak KJ and Ranjana M: Rapid determination of polyphenol, vitamins, organic acids and sugars in *Aegle marmelos* using reverse phase-high performance liquid chromatography. *Journal of Pharmacy Research* 2011; 4 (3):717-719.