

Review Article

Comprehensive Study of Nirgundi Plant: A Survey Report

Rakesh Tiwle^{*1}, D. K. Sanghi¹

¹Shri Laxmanrao Mankar Institute of Pharmacy, Amgoan, Gondia, Maharashtra, India- 441902.

Abstract

The genus *Vitex* contains 270 species distributed around the world. It is an interesting source of potential bioactive molecules, as iridoids compounds, flavonoids, diterpenoids derivatives, phytosteroids, with antioxidant, anti-inflammatory, antimicrobial, Hepatoprotective activity, analgesic and antihistamine. It is a large aromatic shrub distributed throughout India. In ancient times it is used as a female remedy and also for pains in Ayurveda and also in Roman medicine. It is distributed through Indo-Malesia, Europe, Asia, West Indies and India, in India it is found in the outer Himalayas. It is also called as village pharmacy because it is used for the many disease like antibacterial, astringent, febrifuge, vermifuge, sedative, antitumor, tonic various chemical constituents like flavones, glycosides, volatile oil, flavonoids, tri terpenes, and tannins many others were identified in this plant.

Key words: Nirgundi, Antibacterial Activity, Anticancer Activity.

***Corresponding Author: Rakesh Tiwle**, Shri Laxmanrao Mankar Institute of Pharmacy, Amgoan, Gondia, Maharashtra, India- 441902

1. Introduction

Vitex negundo L. commonly known as Nirgundi belongs to family Verbenaceae it is used for the treatment of Analgesic, Antibacterial, Anticonvulsant, dengue, rheumatism, dyspepsia and diarrhea in a folkloric history it is used for cold, cough, asthma and also it strengthens the immune and inhibitory effect on *Xanthomonas axonopodis* and *Pseudomonas solanacearum* and Anticonvulsant activity were examined in the survey report from the local region in the qualitative examination constituents like alkaloids, carbohydrates,

glycosides, phenolic compounds, saponins etc[1]




Figure No. 1 Nirgundi plant (Shri Laxmanrao Mankar Institute of Pharmacy, Amgoan)

Geography Condition

Vitex usually grows from three to nine feet tall; under cultivation it can develop to 20 feet tall. This species is globally distributed in Indo-Malaysia, cultivated in America, Europe, Asia and West Indies. [2] Within India, it is found throughout the greater part of India, ascending to an altitude of 1500m.

Cultivation

It prefers a light well-drained loamy soil.[3] It is widely planted as a hedge plant in between the fields and usually not browsed by the cattle. It produces root suckers which can also be utilized as planting material. An easily grown plant, plants tolerate temperatures down to about 10°C.

Leaf

Leaves are palmately compound, long; 3-5 foliate, petiole 2.5-3.8 cm the middle leaflet is petiolate; in trifoliate leaf, leaflet lanceolate middle leaflet 5-10 cm long and 1.6-3.2 cm broad, with 1-1.3 cm long it consist of β -caryophyllene; sabinene; 4-terpineol; gamma-terpinene; caryophyllene oxide; 1-oceten-3-ol; globulol; 5,3'-dihydroxy-7,8,4'-trimethoxyflavanone; 6-p-hydroxybenzoyl mussaenosidic acid; 2'-p-hydroxybenzoyl mussaenosidic acid viridiflorol; 5,3-dihydroxy,6,7,4 trimethoxyflavanone; 5hydroxy-7,4' dimethoxy flavones; 5,3'-dihydroxy-7,8,4'-trimethoxy flavanone; betulinic acid [3β -hydroxylup-20-(29)-en-28-oic acid]; 5-hydroxy-3,6,7,3',4'-pentamethoxy flavones; 5,7dihydroxy-6,4' dimethoxy flavonone; ursolic acid [2β - hydroxyurs-12-en-28-oic acid]; n-hentriacontanol; β -sitosterol; p-hydroxybenzoic acid; protocatechuic acid;

oleanolic acid; flavonoids angusid; casticin; vitamin-C.[3,4]

Root

Roots are woody, fairly thick, 8-10 cm in diameter; external surface brownish, rough due to the presence of longitudinal fissures and a small rootlets.[5] The bark is very thin and corky portion can be scrapped off easily it consist of chemical constituent like $2\alpha,3\alpha$ -dihydroxyoleana-5,12-dien-28-oic acid; $2\alpha,3\beta$ -diacetoxy-18-hydroxyoleana-5,12- dien-28-oic acid; $2\beta,3\alpha$ -diacetoxyoleana-5,12-dien-28-oic acid; isovitexin, negundin-A; vitexin; (+)-diasyringaresinol; (+)- lyoniresinol; vitrofolal-E; vitrofolal-F, acetyl oleanolic acid; negundin-B; sitosterol; 3-formyl-4,5-dimethyl-8- oxo-5H-6,7-dihydronaphtho (2,3-b)furan.[6]

Bark

It is about 0.3- 0.5 cm thick; rough, lenticelular, outer surface yellowish grey, smooth; fracture short and splintery, longitudinally channeled and transversely cracked; inner surface darker than outer, blackish in colour.[7]

Lamina

shows single layered epidermis having mostly unicellular hairs, bi and multicellular and glandular trichomes being rare; hypodermis 1-3 layered interrupted at places by 4-8 palisade layers containing chlorophyll; [8] a large number of veins enclosed by bundle sheath traverse mesophyll; stomata present only on the ventral surface, covered densely with trichomes; vein-islet of leaf are 23-25 and vein termination number are 5-7 respectively.[9]

Dosage: Nirgundi Juice - 20 to 30 ml per day. Nirgundi leaf Powder - 3 to 6 grams.

Common name of *Vitex negundo* Nirgundi in different languages.[10]

Sanskrit, *Indrani, Nilanirgundi, Nilapushpa, Nirgundi, Nirgundika, Renuka, Sephalika, Shephali, Shvetasurasa, Sindhooka, Sindhuvaram*. **Malyalam** [11] *Bem-nosi, Indrani, Karunocci, Noch-chi, Nochi, Vella-noch-chi* **Marathi** [12] *Nirgunda, Nengar, Nirgur, Lingur, Nirguda, Nirgundi, Limgud, Negumd*. **Hindi** [13] *Mewri, Nengar, Ningori, Nirgandi, Nirgunda, Nisinda, Panikisambhalu, Sambhal, Sambhalu, Nirgundi, Shimalu* **Urdu** [14] *Sambhalu, Tukhm sambhalu*. **Bengali** [15] *Nisinda, Sinduari, Beguna, Nishinda, Nishinde*. **Kannada** [16] *Bilenekki, Bilenekki. Karilakki, Lakkagida, Lakki, Lakki-gida, Lakkili* **Oriya** [17] *Thingkhawilupa, Niligundi* **Tamil** [18] *Nallanocci, Nirkkundi, Nirkundi, Nochi, Sinduvara*.

Essential oil of fresh leaves, flowers and dried fruits[19]

β -guaiene; guaia-3,7-dienecaryophyllene epoxide; ethyl-hexadecenoate; α -selinene; germacren-4-ol; caryophyllene epoxide; (E)-nerolidol; β -selinene; α -cedrene; germacrene D; hexadecanoic acid; p-cymene and valencene. viridiflorol (19.55%), β -caryophyllene (16.59%), sabinene (12.07%), 4-terpineol (9.65%), γ -terpinene (2.21%), caryophyllene oxide (1.75%), 1-oceten-3-ol (1.59%), and globulol (1.05%).[20]

2. Methodology

Preparation of extracts

1. Antibacterial Activity[21]

Leaf of Nirgundi plant dried in the atmosphere followed by it will be crushed [22]. 50g powder was initially soaked in 200 ml of DCM (Dichloro methane) in airtight conical flask in a shaker for 72


hours and then it was filtered through by muslin cloth and then filtered it through Whatman filter paper.[23] The filtrate was collected into airtight brown bottle, similar process was repeated thrice with fresh DCM and the filtrates were pooled together. Followed by the DCM can be removed by using rotary evaporator at low temperature and these dried extract material was stored in the refrigerator.[24] These extract can be sprayed in to the plant important crop plants including potato, tobacco, tomato and groundnut to prevent the pathogenic bacteria, inhibitory effect on *Xanthomonas axonopodis* and *Pseudomonas solanacearum*[25].



Figure No. 2 Antibacterial activity on Tomato plant

2. Anticulscent Activity [26]

Almost all parts of plant are used in the Ayurvedic and Unani system of medicines.[27] *Vitex negundo* (Verbenaceae), is used for inflammatory swelling of joints from acute rheumatism, healing wounds, ulcers and hepatic disorders.[28] Hence it was thought worthwhile to investigate the anti epileptic activity of alcoholic extract of roots of *Vitex negundo*. [29] A survey report of nirgundi plants used in the adivasi areas is shown in the following survey report [30].



MARKETED FORMULATION OF NIRGUNDI PLANTS [31]



Figure No. 3 Marketed formulation of Nirgundi.

3. Conclusion

Ethno botanical and traditional uses of *Vitex negundo* Nirgundi the fact that it is a popular remedy. The above study revealed

the antibacterial activity and anticulscunt activity of *Vitex negundo* and *Allium sativum* against *Pseudomonas solanacearum* and *Xanthomonas axonopodis* extract of flower of *Vitex*

negundo showed higher inhibitory effect. Further, this is the first report that demonstrates the inhibitory effect of *Vitex* the results of this study provide support for the traditional use of *Vitex negundo* as an anticonvulsant drug. Phytochemical screening has shown the presence of alkaloids, carbohydrates, glycosides, phenolic compounds, saponins and sterols in alcoholic extract. The potent activity may be attributed to the presence of these phytoconstituents.

Acknowledgement

The authors would like to acknowledge the assistance provided by kind cooperation of Secretary Shri Keshavrao Mankar Bhavabhuti Shikshan Sanstha Shri Laxmanrao Mankar Institute of Pharmacy Amagoan, Gondia Maharashtra, INDIA.

References

1. Adnaik RS, Pai PT, Mule SN, Naikwade NS, Magdum CS. Laxative Activity of *Vitex negundo* Linn. Leaves. Asian J. Exp. Sci. 2008; 22(1): 159-160.
2. International journal of advances in pharmacy, Biology and chemistry Vol. 1(1), Jan- Mar, 2012, Natural product redience vol 4(3) may-june 2005
3. Liu C, Tseng A, Yang S. Chinese herbal medicine: modern applications of traditional formulas. Florida, USA: CRC Press; 2005.
4. Zheng CJ, Zhao XX, Ai HW, Lin B, Han T, Jiang YP, et al. Therapeutic effects of standardized *Vitex negundo* seeds extract on complete Freund's adjuvant induced arthritis in rats. Phytomedine 2014; 21: 838-846.
5. Vinuchakkaravarthy T, Kumaravel KP, Ravichandran S, Velmurugan D. Active compound from the leaves of *Vitex negundo* L. shows anti-inflammatory activity with evidence of inhibition for secretory phospholipase A2 through molecular docking. Bioinformation 2011; 7: 199-206.
6. Chattopadhyay P, Hazarika S, Dhiman S, Upadhyay A, Pandey A, Karmakar S, et al. *Vitex negundo* inhibits cyclooxygenase-2 inflammatory cytokine-mediated inflammation on carrageenan-induced rat hind paw edema. Pharmacogn Res 2012; 4: 134-137.
7. Baytop T. Therapy with Medicinal Plants (Past and Present) I, Stanbul University Publications, Istanbul, 1984, pp. 252.
8. Honda G, Yes ilada, Tabata M, Sezik E, Fujita T, Takeda Y, Takaishi Y, Tanaka T. Traditional medicine in Turkey VI. Folk medicine in West Anatolia: Afyon, Kutahya, Denizli, Mugla, Aydin provinces. Journal of Ethnopharmacology 1996; 53: 75-87.
9. Gomaa CS, El-Mokhazy MA, Halim FA, El-Sayyad AE. Flavonoids and iridoids from *Vitex agnus-castus*. Planta Medica 1978; 33: 277.
10. Sirait M, Rimpler H, Ha nsel R. Flavonoide aus *Vitex agnus castus* L. Experientia 1962; 18: 72.
11. Wollenweber E, Mann K. Flavonols from fruits of *Vitex agnus-castus*. Planta Medica 1983; 47:126-127.
12. Hoberg E, Meier B, Sticher O. Quantitative high performance liquid chromatographic analysis of casticin in the fruits of *Vitex agnus-castus*. Pharmaceutical Biology 2001; 39: 57-61.
13. Hoberg E, Orjala J, Meier B, Sticher O. Diterpenoids from the fruits of *Vitex agnus-castus*. Phytochemistry 1999; 52: 1555-1558.
14. Sorensen JM, Katsiotis ST. Parameters influencing the yield and composition of the essential oil from Cretan *Vitex agnus-castus* fruits. Planta Medica 2000; 66: 245-250.
15. Saden-Krehula M, Kustrak D, Blazevic N. 4-3-Ketosteroids in flowers and leaves of *Vitex agnus-castus*. PlantaMedica 1990; 56: 547.

16. Alam MI, Gomes A. Snake venom neutralization by Indian medicinal plants (*Vitex negundo* and *Embllica officinalis*) root extracts. Journal of Ethnopharmacology 2003, 86: 75-80.
17. Karunamoorth K, Ramanujam S, Rathinasamy R. Evaluation of leaf extracts of *Vitex negundo* L. against larvae of *Culex tritaeniorhynchus* and repellent activity on adult vector mosquitoes. Parasitol Res. 2008; 103: 545-550.
18. Adnaik RS, Pai PT. Laxative activity of vitex negundo linn. Leaves, Asian J. Exp. Sci. 2008; 22(1): 159-160.
19. Rastogi T, Bhutda V, Moon K, Aswar PB, Khadabadi SS. Comparative Studies on Anthelmintic Activity of *Moringa Oleifera* and *Vitex*. Asian J. Research Che. 2009; 2(2): 217-221
20. Chandramu C, Manohar RD, Krupadanam DGL, Dashavantha RV Isolation characterization and biological activity of betulinic acid and ursolic acid from *Vitex negundo* L. Phytother. Res. 2003; 17:129-134.
21. Chawla AS, Sharma AK, Handa SS. Chemical investigation and anti inflammatory activity of *Vitex negundo* seeds. Part I. Indian J. Chem. 1991; 30B: 773-776.
22. Mukherjee KS, Badruddoza S. Chemical constituents of *Dillenia indica* Linn. And *Vitex negundo* Linn. J. Indian Chem. Soc. 1981; 58:97.
23. Maurya R, Shukla PK, Ashok K. New antifungal flavonoid glycoside from *vitex negundo*. Bioorganic and Medicinal Chemistry 2007; 17:239-242.
24. Gautam LN. Chemical constituents from *vitex negundo* (Linn.) of nepalese origin. Scientific World 2008; 6: 6.
25. Khan MSS, Syeed HS, Haneefuddin M, Akter L, Asmatullah M, Jahan S, et al. Screening and evaluation of antioxidant, antimicrobial, cytotoxic, thrombolytic and membrane stabilizing properties of the methanolic extract and solventsolvent partitioning effect of *Vitex negundo* bark. Asian Pac J Trop Dis 2013; 3: 393-400.
26. Nagarsekar KS, Nagarsenker S, Kulkarni R. Antioxidant and antilipid peroxidation potential of supercritical fluid extract and ethanol extract leaves of *Vitex negundo* Linn. Indian J Pharm Sci 2011; 73: 422-429.
27. Devi PR, Kumari K, Kokilavani C, Effect of *Vitex negundo* leaf extract on the free radicals scavengers in complete freund's adjuvant induced arthritic rats. Indian Journal of Clinical Biochemistry. 2007; 22 (1): 143-147.
28. Tandon VR, Gupta RK, *Vitex negundo* Linn (VN) leaf extract as an adjuvant therapy to standard anti-inflammatory drugs. Indian J Med Res. 2006; 124(4): 447-50
29. Bansod MS, Harle UN, *Vitex negundo* L: Phytochemical constituents, traditional uses and pharmacological properties: Comprehensive review. Pharmacologyonline Newsletter. 2009; 1: 286-302.
30. Jabeen, A., Khan, M., Zafar, M. and Ahmad, F. (2009) 'Indigenous uses of economically important flora of Margallah Hills National Park, Islamabad, Pakistan', African Journal of Biotechnology. 8, 763-784.
31. Vishwanathan, A. S., Basavaraju, R. (2010) 'A Review on vitex negundo L.-A medicinally important plant', EJBS. 3(1), 30-42.